


# Exact Solutions to the Navier-Stokes Equation

Goodarz Ahmadi

Department of Mechanical and Aeronautical Engineering  
Clarkson University  
Potsdam, NY 13699-5727

## Outline

- ▶ Plate Suddenly Set in Motion
- ▶ Oscillating Plate
- ▶ Unsteady Pipe Flows
- ▶ Steady Flows in Noncircular Pipes
- ▶ Elliptic Cross Section Pipes
- ▶ Triangular Cross-Section Pipes



## Similarity Solution

Let

$$t \sim t^1$$

$$y \sim t^a$$

Navier-Stokes

$$1 = 2a$$

$$a = 1/2$$

Similarity Variables

$$\eta = \frac{y}{2\sqrt{\nu t}}$$

$$\frac{u}{U_0} = f(\eta)$$

# Plate Suddenly Set in Motion Clarkson University

**NS**  $\rightarrow$   $f'' + 2\eta f' = 0$

**B.C.**  $\rightarrow$   $f(0) = 1$   
 $f(\infty) = 0$


$\rightarrow$   $f' = ce^{-\eta^2}$

$f = 1 - \frac{2}{\sqrt{\pi}} \int_0^\eta e^{-\eta_1^2} d\eta_1 = 1 - \text{erf}(\eta)$

**Solution**  $\rightarrow$   $u = U_0 \text{erfc}\left(\frac{y}{2\sqrt{vt}}\right)$

ME 637

G. Ahmadi



# Oscillating Plate Clarkson University

$\frac{\partial u}{\partial t} = \nu \frac{\partial^2 u}{\partial y^2}$

**B.C.**  $y = 0 \rightarrow u = U_0 \cos \omega t$   
 $y = \infty \rightarrow u = 0$

**Let**  $u = U_0 e^{-ky} \cos(\omega t - ay)$

ME 637

G. Ahmadi

# Oscillating Plate Clarkson University

$\frac{\partial u}{\partial t} = -\omega U_0 e^{-ky} \sin(\omega t - ay)$        $\frac{\partial u}{\partial y} = U_0 e^{-ky} (-k \cos(\omega t - ay) + a \sin(\omega t - ay))$

**Navier-Stokes Equation**

$-\omega \sin \theta = \nu((k^2 - a^2) \cos \theta - 2ak \sin \theta)$

**Matching**  $\rightarrow$   $a^2 = k^2$        $\omega = 2ak\nu = 2k^2\nu$

**Solution**  $u = U_0 e^{-ky} \cos(\omega t - ky)$        $k = \sqrt{\frac{\omega}{2\nu}}$


ME 637

G. Ahmadi

# Unsteady Flow in a Tube Clarkson University

**Navier Stokes**

$$\frac{\partial v_z}{\partial t} = -\frac{1}{\rho} \frac{dP}{dz} + \nu \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial v_z}{\partial r} \right)$$



ME 637

G. Ahmadi

# Unsteady Flow in a Tube Clarkson University

**Nondimensional Variables**

$$\xi = \frac{r}{R}$$

$$\tau = \frac{\mu t}{\rho R^2} = \frac{\nu t}{R^2}$$

$$v_z = -\frac{1}{4\mu} \frac{dP}{dz} R^2 \phi(\xi)$$

**Navier-Stokes**

$$\frac{\partial \phi}{\partial \tau} = 4 + \frac{1}{\xi} \frac{\partial}{\partial \xi} \left( \xi \frac{\partial \phi}{\partial \xi} \right)$$

ME 637

G. Ahmadi

# Unsteady Flow in a Tube Clarkson University

**Boundary Conditions**

$$\xi = 1$$

$$\phi = 0$$

$$\tau = 0$$

$$\phi = 0$$

**Changing Variable**

$$\phi = 1 - \xi^2 - \psi$$

ME 637

G. Ahmadi

# Unsteady Flow in a Tube Clarkson University

**Navier Stokes**

$$\frac{\partial \psi}{\partial \tau} = \frac{1}{\xi} \frac{\partial}{\partial \xi} \left( \xi \frac{\partial \psi}{\partial \xi} \right)$$

**Boundary Conditions**

$$\xi = 1$$

$$\psi = 0$$

$$\tau = 0$$

$$\psi = 1 - \xi^2$$

ME 637

G. Ahmadi

# Unsteady Flow in a Tube Clarkson University

## Separation of Variables

$$\psi = F(\xi)T(\tau)$$

$$\frac{\dot{T}}{T} = \frac{1}{F\xi} \frac{d}{d\xi} \left( \xi \frac{dF}{d\xi} \right) = -\alpha^2$$

$$\dot{T} + \alpha^2 T = 0$$

$$T = Ce^{-\alpha^2 \tau}$$

## Bessel Equation

$$\xi^2 \frac{d^2 F}{d\xi^2} + \xi \frac{dF}{d\xi} + \alpha^2 \xi^2 F = 0$$

## Bessel Functions

$$F = AJ_0(\alpha\xi) + BY_0(\alpha\xi)$$

ME 637

G. Ahmadi

# Unsteady Flow in a Tube Clarkson University

## Boundary Conditions

$$Y_0(0) \rightarrow \infty$$

$$F(0) \sim \text{finite} \Rightarrow B = 0$$

$$F(1) = 0 \Rightarrow J_0(\alpha) = 0$$

## Eigenvalues

$$\alpha_1 = 2.405$$

$$\alpha_2 = 5.52$$

$$\alpha_3 = 8.654$$

## General Solution

$$\psi = \sum_n A_n e^{-\alpha_n^2 \tau} J_0(\alpha_n \xi)$$

ME 637

G. Ahmadi

# Unsteady Flow in a Tube Clarkson University

## Initial Condition

$$1 - \xi^2 = \sum_n A_n J_0(\alpha_n \xi)$$

$$A_n = \frac{\int_0^1 (1 - \xi^2) \xi J_0(\alpha_n \xi) d\xi}{\int_0^1 \xi J_0^2(\alpha_n \xi) d\xi} = \frac{8}{\alpha_n^3 J_1^2(\alpha_n)}$$

## Solution

$$\psi = 8 \sum_n \frac{e^{-\alpha_n^2 \tau} J_0(\alpha_n \xi)}{\alpha_n^3 J_1^2(\alpha_n)}$$


ME 637

G. Ahmadi

# Unsteady Flow in a Tube Clarkson University

## Solution

$$\varphi = 1 - \xi^2 - 8 \sum_n \frac{J_0(\alpha_n \xi)}{\alpha_n^3 J_1^2(\alpha_n)} e^{-\alpha_n^2 \tau}$$




ME 637

G. Ahmadi


# Noncircular Pipe Flows Clarkson University

**Navier Stokes**



$$\nabla^2 W = \frac{1}{\mu} \frac{dP}{dz} = \text{const}$$

$$W = 0$$



ME 637

G. Ahmadi


# Elliptical Pipes Clarkson University

**Ellipse**

$$\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1$$



$$w = A \left(1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}\right)$$



ME 637

G. Ahmadi

# Elliptical Pipes Clarkson University

**NS**

$$\nabla^2 w = -A \left(\frac{2}{a^2} + \frac{2}{b^2}\right) = -\frac{2A(a^2 + b^2)}{a^2 b^2} = \frac{1}{\mu} \frac{dP}{dz}$$

$$A = -\frac{1}{2\mu} \frac{dP}{dz} \frac{a^2 b^2}{a^2 + b^2} \left(1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}\right)$$

**Flow Rate**

$$Q = \iint w dx dy$$

$$Q = -\frac{\pi}{4\mu} \frac{dP}{dz} \frac{a^3 b^3}{a^2 + b^2}$$

ME 637


G. Ahmadi

# Triangular Pipes Clarkson University

$$f(x, y) = (x - a)(x - \sqrt{3}y + 2a)(x + \sqrt{3}y + 2a) = 0$$

$$(x - \sqrt{3}y + 2a) = 0$$

$$(x + \sqrt{3}y + 2a) = 0$$



ME 637

G. Ahmadi

# Triangular Pipes Clarkson University

**Let**  $w = Af(x, y)$

**NS**  $\nabla^2 w = A\nabla^2 f(x, y) = 12aA = \frac{1}{\mu} \frac{dP}{dz}$

**Solution**

$$A = \frac{1}{12\mu a} \frac{dP}{dx}$$

$$w = \frac{1}{12\mu a} \frac{dP}{dx} (x-a)(x-\sqrt{3}y+2a)(x+\sqrt{3}y+2a)$$

ME 637

G. Ahmadi

# Concluding Remarks Clarkson University

- **Plate Suddenly Set in Motion**
- **Oscillating Plate**
- **Unsteady Pipe Flows**
- **Steady Flows in Noncircular Pipes**
- **Elliptic Cross Section Pipes**
- **Triangular Cross-Section Pipes**

ME 637

G. Ahmadi

Clarkson University

# Thank you!

# Questions?

ME 637

G. Ahmadi